REFERENCES:

1. Ali M and Knight A. Comparative healthcare: Diabetes Mellitus. Australas Med J.2009;1(5):1-9.
2. American Diabetes Association. Diagnosis and Classification of Diabetes Mellitus . *Diabetes Care*. 2009;32(Suppl 1):S62-S67. doi:10.2337/dc09-S062
3. Ann Mary Swaroop, Christina Varghese, Josmine Jose, Maheswari E., Pramila Kalra.*Impact of patient counseling on knowledge , attitude practice and medication adherence in Type-2Diabetes Mellitus* European Journal of Pharmaceutical and Medical Reserch,2016,3(4), 231-235
4. Authoritative Institute of Health and Welfare (AIHW). National health priority areas. 2013.http://www.aihw.gov.au/national-health-priority-areas
5. Beardsley RS. Review of literature: oral patient counseling by pharmacists. Proceedings of the National Symposium on oral counseling by pharmacists about prescription medicines. 1997 Sep 19-21; Lansdowne, Virginia
6. Butt M, Mhd Ali A, Bakry MM, Mustafa N. Impact of a pharmacist led diabetes mellitus intervention on HbA1c, medication adherence and quality of life: A randomised controlled study. *Saudi Pharmaceutical Journal : SPJ*. 2016;24(1):40-48. doi:10.1016/j.jsps.2015.02.023
7. Buysschaert M, Hermans MP. Non-pharmacological management of type 2 diabetes. Acta Clin Belg 2004;59:14-9.
8. FIP.org. Global pharmacy workforce and migration report: a call for action. 2016. Availablefrom:https://www.fip.org/files/fip/publications/ PharmacyWorkforceMigration.pdf. Accessed October 28, 2016.
9. Haynes, R.B., McDonald, H., Garg,A.X., & Montgue,P.Interventions for helping patients to follow prescriptions for medications.Journal of the American Medical Association 2002; Vol 288, Page No.2880-2883
10. Heidi M, Harper A and Berger BA et al. Pharmacists predisposition to communicate-Desire to counsel and job satisfaction. Am J Pharm Educ 1992; 56: 252- 8
11. Iglay K, Cartier SE, Rosen VM, et al. Meta-analysis of studies examining medication adherence, persistence, and discontinuation of oral antihyperglycemic agents in type 2diabetes.Curr Med Res Opin. 2015;31(7):1283–1296.
12. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. AMJ 2014, 7, 1,45-48. http//dx.doi.org/10.4066/AMJ.2014.1979)
13. Kennedy-Martin T, Boye KS, Peng X. Cost of medication adherence and persistence in type 2 diabetes mellitus: a literature review. Patient Prefer Adherence. 2017 Jun

30;11:1103-1117

14. Lee VW, Pang KK, Hui KC et al. Medication adherence: is it a hidden drug-related problem in hidden elderly? Geriatric Geritol Int 2013; 13: 978–985
15. Malathy R, Narmadha MP, Ramesh S, Alvin JM, Dinesh BN. Effect of a diabetes counseling programme on KAP among diabetic patients in Erode district of South India.J young Pharm 2011; 3(1): 65-72.
16. Mathew E, Ahmed M, Hamid S, Abdulla F, Batool K. Hypertension and dyslipidaemia in Type 2 diabetes mellitus in United Arab Emirates. Australas Med J. 2010; 3(11):699- 706.
17. Morisky, D.E, Green, L.W.,& Levine, D.M.Concurrent and predictive vlidity of a self-reported measure of medication adherence.Medical Care 2000 Vol 24, Page No.67-74.
18. National Diabetes Education Program. Centers for Disease Control and Prevention.
19. National service frameworks and strategies. National Health Services. July 2011. Accessed on 13 Dec. 2013)
20. Piette,J.D, Heisler, M., Ganoz, D., McCarthy, J.F., & Valenstein, M. Differential medication adherence among patients with diabetes and hypertension.2007; Vol. 58,Page No.207-212
21. prevalence of diabetes for 2011 and 2030. Diabetes Res Clin Pract. 2011;94:311-21)
22. Schoenberg NE. The relationship between perceptions of social suppert and adherence to dietary recommendations among African-American elders with hypertension.Int J Aging HUM Dev Page No.279-297
23. State-based diabetes prevention and control program. Centers for Disease Control and Prevention
24. Stoehr GP, Lu SY, Lavery L et al. Factors associated with adherence to medication regimens in older primary care patients: the Steel Valley Seniors Survey. Am J Geriatr Pharmacother 2008; 6: 255–263. 50
25. The Seventh Report of the Joint National Committee on Prevention,Detection,Evaluation and Treatment of High Blood Pressure.The JNC 7 Report. JAMA 2003,Page No.2560-2572
26. ThiruchselvamT, NaglieG,MoineddinRet al.Riskfactors formedicationnonadherenceinolderadultswithcognitive impairment who live alone. Int J Geriatr Psychiatry 2012; 27: 1275–1282. 51
27. USP medication counseling behaviour guideline. USP DI update volumes I and II Rockville
28. Venkatesan R, Devi ASM, Parasuraman S, Sriram S. Role of community pharmacists

in improving knowledge and glycemic control of type 2 diabetes. *Perspectives in Clinical Research*. 2012

29. Whiting Dr, Guariguata L, Weil C, Shawj. IDF Diabetes atlas: Global estimates of the

30. WHO World Health Organization, Global Report on Diabetes-2016

31. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes-estimates for the year 2000 and projections for 2030. Diabetes Care. 2004;27(3):1047-53

32. Wilper AP,Woolhandler S, Lasser KE,McCormick D, Bor DH, Himmelstein DU. A national study of chronic disease prevalence and access to care in uninsured US adults. An Intern Med.2008;149(3),Page No.170-17